



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

for December, 1897, gives additional corroboration of the former claim, since he shows that the secretion digests fibrin in the presence of one per cent. hydrocyanic acid, and that its enzyme retains its digestive activity when kept for several weeks in pure glycerine. His studies do not reach to the secretion of the necessary acid, which, in one species at least, is present in the liquid of unopened pitchers, and therefore is not the result of stimulation by the presence of foreign bodies.

Primitive Angiosperms.—From a morphological study of *Naias* and *Zannichellia*,¹ Professor Campbell shows that both anthers and ovules are axial structures, approaching, as he believes, more closely to the sporangia of Pteridophytes than do those of any other angiosperm, and he seems inclined to look upon these genera as standing nearer to the diverging point of Isoetaceæ and monocotyledons than do most representatives of the latter group.

New Species of Pectis.—Mr. M. L. Fernald, of the Gray Herbarium, contributes to the knowledge of Mexican plants by publishing in vol. xxiii, no. 5, of the *Proceedings of the American Academy of Arts and Sciences* a paper on some rare and undescribed plants collected at Acapulco by Dr. Edward Palmer in 1894, and a systematic study of the genus *Pectis*, including species of the United States as well as Mexico. Of this genus, *P. Lessingii*, *P. prostrata*, var. *cylindrica*, and var. *urceolata*, *P. sinalensis*, *P. depressa*, *P. capillaris*, var. *paucicapitata*, *P. filipes*, var. *subnuda*, *P. Pringlei*, *P. Rosei*, *P. elongata*, var. *Schottii*, *P. ambigua*, and *P. linifolia*, var. *marginalis* are described as new.

Botanical Notes.—Botanists will be interested in knowing that the herbarium and notes of the late M. S. Bebb, a collection invaluable for any systematic study of North American willows, have been purchased by the Field Columbian Museum of Chicago.

An article on "The North American Genus *Sarracenia*," illustrated by a reproduction of a photograph of *S. Chelsoni* (*S. rubra* × *purpurea*), is to be found in *Gartenwelt*, of Berlin, of Dec. 26, 1897.

Students of European botany, who have found difficulty in resigning themselves to the use of one name for the terra-cotta-flowered and blue-flowered forms of the poor man's weather-glass, which they

¹ D. H. Campbell, Contributions to Biology from the Hopkins Seaside Laboratory of the Leland Stanford Junior University. XI, A Morphological Study of *Naias* and *Zannichellia*. Reprinted from *Proceedings of the California Academy of Sciences*, 3d ser., Bot., vol. i. San Francisco, Cal., 1897. 61 pp., 5 pl.